

Optimal PMU Placement Problem in Octahedral Networks

S. Prabhu^{a,*}, S. Deepa^b, Rajvikram Madurai Elavarasan^c, Eklas Hossain^d

^aDepartment of Mathematics, Rajalakshmi Engineering College, Chennai 602105, India

^bDepartment of Mathematics, Easwari Engineering College, Chennai 600089, India

^cClean and Resilient Energy Systems Laboratory, Texas A&M University, Galveston, TX 77553, USA

^dOregon Renewable Energy Center, Department of Electrical Engineering and Renewable Energy
Oregon Institute of Technology, Klamath Falls, OR 97601, USA

Abstract

Power utilities must track their power networks to respond to changing demand and availability conditions to ensure effective and efficient operation. As a result, several power companies employ phase measuring units (PMUs) to check their power networks continuously. Supervising an electric power system with the fewest possible measurement equipment is precisely the vertex covering graph-theoretic problem, in which a set D is defined as a power dominating set (PDS) of a graph if it supervises every components (vertices and edges) in the system (with a couple of rules). The $\gamma_p(G)$ is the minimal cardinality of a PDS of a graph G . In this present study, the PDS is identified for octahedral networks.

Mathematics Subject Classification 2020: 05C69 · 90C27

Key Words and Phrases: Dominating set; Phase measurement unit; Private neighbor; Power domination; Octahedral network

1 Introduction

A power network is made up of electrical hubs and transmitting cables that connect them. Electric power companies must constantly track the condition of their systems. The magnitude of the voltage at loads and the system process at generators must both be monitored. The placement of PMUs

*Corresponding author : drsavariprabhu@gmail.com

at certain locations within the device must be controlled. Due to the increasing price of PMUs, it's critical to employ as little as possible while still being able to track the entire system. This Power system observability with minimal PMU placement is a problem that Haynes et al. [19] introduced as a theoretical graph problem and dubbed the power dominating set (PDS) problem after it was demonstrated in [3].

Let G be a connected network with vertex set $V(G)$ and edge set $E(G)$. For any $v \in V(G)$ an open r -neighborhood of v is $N_r(v) = \{u \in V(G) : d_G(u, v) = r\}$, where $d_G(u, v)$ is the distance (number of edges in u - v path) between the u to v . Similarly $N_r[v] = N_r(v) \cup \{v\}$. The degree of v in $V(G)$ is $d_G(v) = |N_1(v)|$. Denote $[\alpha] := \{1, 2, \dots, \alpha\}$ and $[\alpha]_0 := \{0, 1, 2, \dots, \alpha\}$. Also to denote $\{\alpha + 1, \alpha + 2, \dots, 2\alpha + 1\}$ we use the notation $[2\alpha + 1] - [\alpha]$.

A dominant subset is a proper subset D of $V(G)$ if any node in $V(G)$ that is not in D must have at least one adjacent node in D . The least cardinality of all possible D is $\gamma(G)$. If all nodes of $V(G)$ can be recursively observed by either domination or propagation, a subset D is called PDS (power dominating set).

1. **Rule (i): (Domination)**

$$M(D) \leftarrow D \cup N(D)$$

2. **Rule (ii): (Propagation)**

$$\exists, v \in M(D) \text{ s.t. } (V(G) - M(D)) \cap N(v) = \{w\}$$

$$M(D) \leftarrow M(D) \cup \{w\}$$

A subset D is named as PDS for G with $M(D) = V(G)$. The least positive integer representing a PDS is $\gamma_p(G)$.

The problem of finding PDS is NP-complete in general. Even for graph classes like chordal, split and bipartite, it remains NP-complete [19]. In [1, 18, 19], various algorithms for computing the PDS for a specific class of graphs were described. An improved algorithms with complexity results were reported in [18]. This problem is studied for block graphs [34], circular-arc graphs [26], hypercubes [8, 11], grids [16], generalized Petersen graphs [4, 9, 22, 33], permutation graphs [32], planar graphs with small diameter [37], maximal planar graphs [12], Knodel graphs and Hanoi graphs [20], de Bruijn graphs and Kautz graphs [24], regular claw-free graphs [28], and certain chemical graphs [30]. This problem is also discussed for Cartesian product of graphs in [4, 23], tensor and strong product in [15], corona product and join of graphs in [35], and for some other graph products were discussed in [5]. The lower bounds for this problem is discussed in [17]. An

upper bound for one component graph with $n > 4$ is presented, and few extremal graphs concerning PDS are characterized in [36]. The Nordhaus-Gaddum results of this problem were reported in [5].

Straight forward generalization of PDS problem is done in [9] as the k -PDS problem. It is trivial to note that when $k = 1$, this problem converges to the original PDS problem and when $k = 0$ it is a traditional domination problem. This k -PDS problem is discussed for Sierpiński graphs [14], block graphs [31], regular graphs [13], certain interconnection networks [29], and weighted trees [10]. In [7], the complexity of power dominating throttling is discussed. The infectious power domination is introduced in [6], along with a general bound for determining the influence of particular hypergraph operations.

2 Octahedral and its Derived Networks




Figure 1: (a) Unit octahedron; (b) Twin octahedron

The octahedral structures are introduced in [2]. A platonic solid's corresponds to a polyhedral graph, which is called an octahedron graph. The unit octahedron contains 6 vertices and 12 edges. We define a twin octahedron in $OH(n)$ as two octahedrons sharing exactly one common vertex. The structural graph of the unit octahedron and twin octahedron and the vertex representations are depicted in Figure 1. For more information on these graphs, it can be seen from [21, 25]. Different ways of connecting unit octahedron derive the varieties of octahedral structures.

2.1 Octahedral Network

An n -dimensional octahedral network has $27n^2 + 3n$ vertices and $72n^2$ edges. It is noted that $OH(n)$ has $6n^2$ unit octahedron and $3n^2$ edge-disjoint twin octahedron. Figure 2(a) gives an idea about the extension of $OH(n)$ network and its addressing scheme is depicted in Figure 2(b). $V(OH(n)) =$





Figure 2: (a) Construction of $OH(2)$; (b) Addressing scheme of $OH(2)$


$\bigcup_{i=1}^2 P_i \cup \bigcup_{i=1}^2 Q_i$ where, $P_1 = \{p_{i,j} : i \in [n+1] \ \& \ j \in [n+i-1]\}$, $P_2 = \{p_{i,j} : i \in [2n+1] - [n+1] \ \& \ j \in [3n-i+1]\}$, $Q_1 = \{q_{i,j} : i \in [n] \ \& \ j \in [2(n+i)]\}$, and $Q_2 = \{q_{i,j} : i \in [2n] - [n] \ \& \ j \in [2(3n-i+1)]\}$.

2.2 Dominated Octahedral Network

The structural graph of n dimensional dominated octahedral network $DOH(n)$, $n \geq 2$ and its addressing scheme is portrayed in Figure 3. It has $81n^2 - 75n + 24$ vertices and $216n^2 - 216n + 72$ edges. It contains $18n^2 - 18n + 6$ unit octahedron and hence $9n^2 - 9n + 3$ twin octahedron. The $V(DOH(n)) = \bigcup_{i=1}^3 P_i \cup \bigcup_{i=1}^3 Q_i$, where $P_1 = \{p_{i,j} : i \in [n] \ \& \ j \in [3i-2]\}$, $P_2 = \{p_{i,j} : i \in [3n-1] - [n] \ \& \ j \in [\frac{((-1)^{i-n-1} - 1)}{2} + 3n-1]\}$, $P_3 = \{p_{i,j} : i \in [4n-1] - [3n-1] \ \& \ j \in [12n-3i-2]\}$, $Q_1 = \{q_{i,j} : i \in [n-1] \ \& \ j \in [6i-2]\}$, $Q_2 = \{q_{i,j} : i \in [3n-1] - [n-1] \ \& \ j \in [6n-2]\}$, and $Q_3 = \{q_{i,j} : i \in [4n-2] - [3n-1] \ \& \ j \in [24n-6i-8]\}$.



(a)



(b)

Figure 3: (a) Construction of $DOH(2)$; (b) Addressing Scheme of $DOH(2)$

2.3 Rectangular Octahedral Network of Type I and Type II




Figure 4: (a) $ROH^1(3, 9)$; (b) $ROH^1(3, 8)$; (c) $ROH^1(4, 9)$; (d) $ROH^1(4, 8)$.

In line with [27], we introduce the rectangular octahedral network in this subsection. The rectangular octahedral network of Type I denoted by $ROH^1(m, n)$ is derived by arranging octahedron in a two-dimensional plane so that the first octahedron whose apex is facing down. Rectangular octahedral network of Type II denoted by $ROH^2(m, n)$ is derived by presenting octahedrons in a two-dimensional plane similar to $ROH^1(m, n)$ with the condition that the first octahedron, which is the north west corner most unit octahedron whose apex should be facing the top. Rectangular octahedral network of Type I has $\frac{1}{2}(9mn + 2m + n - 1)$ vertices and Type II has $\frac{1}{2}(9mn + 2m + n + 1)$ vertices for m even and n odd. For other cases, both Type I and Type II has $\frac{1}{2}(9mn + 2m + n)$ vertices. Both types of these structures have $12mn$ edges. Different cases of $ROH^1(m, n)$ and their addressing schemes are portrayed in Figure 4. It is clear that $ROH^1(m, n) \cong ROH^2(m, n)$ for all m and n except m even and n odd. The non-isomorphic case is




Figure 5: $ROH^2(4, 7)$

depicted in Figure 5. The vertex set of Type I rectangular octahedral network $V(ROH^1(m, n)) = P \cup Q$, where, $P = \{p_{i,j} : i \in [m+1] \ \& \ j \in [\frac{n}{2}]\}$, $Q = \{q_{i,j} : i \in [m] \ \& \ j \in [n+1]\}$, for n even and $P = \{p_{i,j} : i \in [m+1] \ \& \ j \in [\frac{n+(-1)^i}{2}]\}$, $Q = \{q_{i,j} : i \in [m] \ \& \ j \in [n+1]\}$. for n odd.

The vertex set $V(ROH^2(m, n)) = P \cup Q$, where $P = \{p_{i,j} : i \in [m+1] \ \& \ j \in [\frac{n-(-1)^i}{2}]\}$ and $Q = \{q_{i,j} : i \in [m] \ \& \ j \in [n+1]\}$ for non-isomorphic case of Type II rectangular octahedral network.

3 Main Results

Theorem 1. *Let G be any octahedral structure, H be the edge disjoint subgraph of G isomorphic to twin octahedron and D be the PDS of G . Then $V(H) \cap D \neq \emptyset$.*

Proof. Since H is an edge-disjoint subgraph of G , the nodes a, b, c, d are the only nodes through which the rest of the graph is connected to H . See Figure 6. Suppose the contrary that there exists an H such that $D \cap V(H) = \emptyset$.

Case 1: $N(D) \cap V(H) = \emptyset$

Here $\{a, b, c, d\}$ are not dominated. Suppose that they are observed by propagation, then the vertices $\{x, x', x'', a', b', c', d'\}$ cannot be observed by propagation as at least three vertices of this set are adjacent to each vertex in $\{a, b, c, d\}$. This is the reason for failure of propagation at $\{a, b, c, d\}$. Therefore D must contain some vertices of H . A contradiction.




Figure 6: Edge disjoint subgraph H of G

Case 2: $N(D) \cap V(H) \neq \emptyset$

Here $\{a, b, c, d\}$ are dominated. But due to the fact that $d_G(u) \geq 4$, for $u \in \{a, b, c, d\}$, the propagation at u fails. Therefore the vertices $\{x, x', x'', a', b', c', d'\}$ are not observed. This contradicts the assumption of D . \square


3.1 Power Domination in Octahedral Structures

Theorem 2. For $n \geq 1$, $\gamma_p(OH(n)) = 3n^2$.


Proof. Let D be the PDS of $OH(n)$. Since there are $3n^2$ copies of twin octahedron in $OH(n)$. By Theorem 1, D must have at least $3n^2$ vertices. Therefore $|D| \geq 3n^2$. The reverse inequality is clear from the following choices.

Case 1: n odd.

For this case the PDS $D = \bigcup_{i=1}^2 D_i^P \cup \bigcup_{i=1}^2 D_i^Q$, where $D_1^P = \{p_{i,n+j} : i \in [n+1] - \{1\} \ \& \ j \in [i-1]\}$, $D_2^P = \{p_{n+i+1,n+j} : i \in [n-1] \ \& \ j \in [n-i]\}$, $D_1^Q = \{q_{i,2j} : i \in [n] \ \& \ j \in [n]\}$, and $D_2^Q = \{q_{n+i,2j} : i \in [n] \ \& \ j \in [n]\}$. See Figure 7(a).



(a)



(b)

Figure 7: Power dominating set in octahedral network (a) $OH(3)$; (b) $OH(2)$

Case 2: n even.

For this the PDS is $D = \bigcup_{i=1}^2 D_i^P \cup \bigcup_{i=1}^2 D_i^Q$, where $D_1^P = \{p_{i,j+1} : i \in [n+1] - \{1\} \ \& \ j \in [i-1]\}$, $D_2^P = \{p_{n+i+1,j+1} : i \in [n-1] \ \& \ j \in [n-i]\}$, $D_1^Q = \{q_{i,2j}, q_{i,n+2i+2j-1} : i \in [n] \ \& \ j \in [\frac{n}{2}]\}$, and $D_2^Q = \{q_{n+i,2j}, q_{n+i,4n-2i-2j+3} : i \in [n] \ \& \ j \in [\frac{n}{2}]\}$. See Figure 7(b). □

Theorem 3. For $n \geq 1$, $\gamma_p(DOH(n)) = 9n^2 - 9n + 3$.

Proof. Let $D \subseteq V(DOH(n))$ be the PDS.

There are $9n^2 - 9n + 3$ copies of twin octahedron in $DOH(n)$. By Theorem 1, every twin octahedron must contribute one vertex to D . Hence $|D| \geq 9n^2 - 9n + 3$.

Now let us prove the reverse inequality by exhibiting the PDS of cardinality $9n^2 - 9n + 3$.




Figure 8: Power dominating set in dominated octahedral network $DOH(2)$.

The PDS of dominated octahedral network $DOH(2)$ is depicted in Figure 8. The set $D = \bigcup_{i=1}^3 D_i^P \cup \bigcup_{i=1}^3 D_i^Q$ exhibits the PDS for $DOH(n)$, where $D_1^P = \{p_{i,3j-1} : i \in [n] - \{1\} \ \& \ j \in [i-1]\}$, $D_2^P = \{p_{n+2i-1,3j-2} : i \in [n] \ \& \ j \in [n]\} \cup \{p_{n+2i,3j-1} : i \in [n-1] \ \& \ j \in [n-1]\}$, $D_3^P = \{p_{3n+i-1,3j-1} : i \in [n-1] \ \& \ j \in [n-i]\}$, $D_1^Q = \{q_{i,3j} : i \in [n-1] \ \& \ j \in [2i-1]\}$, $D_2^Q = \{q_{i,3j} : i \in [3n-1] - [n-1] \ \& \ j \in [2n-1]\}$, and $D_3^Q = \{q_{i,3j} : i \in [4n-2] - [3n-1] \ \& \ j \in [8n-2i-3]\}$. \square

Theorem 4. Let G be a rectangular octahedron network $ROH^1(m, n)$, $m, n \geq 1$. Then

$$\gamma_p(G) = \begin{cases} \frac{mn+1}{2} & : m \text{ odd and } n \text{ odd} \\ \frac{mn}{2} & : \text{otherwise} \end{cases}$$

Proof. The proof is arranged as follows.

Case 1: n odd.

Let $V(G) = \{p_{i,j} : i \in [m+1], j \in [\frac{n+(-1)^i}{2}]\} \cup \{q_{i,j} : i \in [m], j \in [n+1]\}$.

Case 1.1: m is odd.

Here we have $\frac{mn-1}{2}$ edge-disjoint twin octahedron and one edge-disjoint unit octahedron. By Theorem 1, we have $\gamma_p(G) \geq \frac{mn-1}{2} + 1$. To prove the reverse inequality by exhibiting PDS in G . The following set $D = D^P \cup D^Q \cup \{p_{m+1, \frac{n+1}{2}}\}$ exhibits the PDS, where $D^P = \{p_{2i,j} : i \in [\frac{m-1}{2}], j = \frac{n+1}{2}\}$ and $D^Q = \{q_{i,2j} : i \in [m], j \in [\frac{n-1}{2}]\}$.

Case 1.2: m is even.

Since m is even, there are $\frac{mn}{2}$ twin octahedron. By Theorem 1, we have $\gamma_p(G) \geq \frac{mn}{2}$. The PDS $D = \{p_{2i,j} : i \in [\frac{m}{2}], j = \frac{n+1}{2}\} \cup \{q_{i,2j} : i \in [m], j \in [\frac{n-1}{2}]\}$ is an evidence for the reverse inequality $\gamma_p(G) \leq \frac{mn}{2}$.

Case 2: n even.

Let $V(G) = \{p_{i,j} : i \in [m+1], j \in [\frac{n}{2}]\} \cup \{q_{i,j} : i \in [m], j \in [n+1]\}$ with cardinality $\frac{1}{2}(9mn + 2m + n)$ and $12mn$ edges. The proof is similar to the rest of the choice of values of m and n along with the PDS $D = \{q_{i,2j} : i \in [m], j \in [\frac{n}{2}]\}$.

□

Theorem 5. Let G be a rectangular octahedron network $ROH^2(m, n)$, $m, n \geq 1$ with m even and n odd. Then $\gamma_p(G) = \frac{mn+2}{2}$.

Proof. Let $V(G) = \{p_{i,j} : i \in [m+1], j \in [\frac{n-(-1)^i}{2}]\} \cup \{q_{i,j} : i \in [m], j \in [n+1]\}$.

Since n is odd, there are $\frac{n-1}{2}$ twin octahedrons in the first and last row of the $V(G)$ and one unit octahedron in each of these rows. Hence there are $\frac{mn-2}{2}$ twin octahedrons in $ROH^2(m, n)$ say $TO_1, TO_2, \dots, TO_{\frac{mn-2}{2}}$. The $V(G) \setminus \bigcup_{i=1}^{\frac{mn-2}{2}} V(TO_i) = S_1 \cup S_2$, where $S_1, S_2 \in \{\{N[p_{1,1}] \cup N[q_{1,1}] \setminus \{q_{1,2}\}\}, \{N[p_{1, \frac{n+1}{2}}] \cup N[q_{1, n+1}] \setminus \{q_{1, n}\}\}, \{N[p_{m+1, 1}] \cup N[q_{m, 1}] \setminus \{q_{m, 2}\}\}, \{N[p_{m+1, \frac{n+1}{2}}] \cup N[q_{m, n+1}] \setminus \{q_{m, n}\}\}\}$. Assume $D = \{p_{2i+1, j} : i \in [\frac{m}{2} - 1] \ \& \ j = \frac{n+1}{2}\} \cup \{q_{i, 2j} : i \in [m] \ \& \ j \in [\frac{n-1}{2}]\}$, such that $M[D] = V(G) \setminus \{S_1 \cup S_2\}$. Since $S_1 \cap S_2 = \emptyset$, we have $\gamma_p(G) = |D| + 2$. Which results $\gamma_p(G) \leq \frac{mn+2}{2}$.

To prove the reverse inequality $\gamma_p(G) \geq \frac{mn+2}{2}$, assume the contrary that $\gamma_p(G) = \frac{mn}{2}$. It is clear by structure, there are $\frac{mn}{2} - 1$ twin octahedrons. By Theorem 1, any PDS D must contain $\frac{mn}{2} - 1$ vertices. The remaining one vertex is not sufficient to power dominate the vertices of S_1 and S_2 due to the fact that $S_1 \cap S_2 = \emptyset$, a contradiction. □

4 Conclusion

This investigation offers exhaustive work on the power domination problem of various networks. It also discusses the optimal PMU placement problem in octahedral networks. The lower bound is attained using edge-disjoint subgraph-technique, and the upper bound is obtained by exhibiting the PDS in an octahedral network. Using these techniques, the tasks mentioned above of placing optimal PMU placement problem with more accurate, optimal and reliable. The problem of placing optimal PMUs for other networks are under investigation.

References

- [1] Ashkan Aazami and Kael Stulp. Approximation algorithms and hardness for domination with propagation. *SIAM Journal on Discrete Mathematics*, 23:1382–1399, 1 2009.
- [2] Micheal Arockiaraj, S. Ruth Julie Kavitha, Krishnan Balasubramanian, and Jia-Bao Liu. On certain topological indices of octahedral and icosahedral networks. *IET Control Theory Applications*, 12:215–220, 1 2018.
- [3] T.L. Baldwin, L. Mili, M.B. Boisen, and R. Adapa. Power system observability with minimal phasor measurement placement. *IEEE Transactions on Power Systems*, 8:707–715, 5 1993.
- [4] Roberto Barrera and Daniela Ferrero. Power domination in cylinders, tori, and generalized petersen graphs. *Networks*, 58:43–49, 8 2011.
- [5] Katherine F. Benson, Daniela Ferrero, Mary Flagg, Veronika Furst, Leslie Hogben, and Violeta Vasilevska. Nordhaus–gaddum problems for power domination. *Discrete Applied Mathematics*, 251:103–113, 12 2018.
- [6] Beth Bjorkman. Infectious power domination of hypergraphs. *Discrete Mathematics*, 343:111724, 3 2020.

- [7] Boris Brimkov, Joshua Carlson, Illya V. Hicks, Rutvik Patel, and Logan Smith. Power domination throttling. *Theoretical Computer Science*, 795:142–153, 11 2019.
- [8] Dennis J. Brueni and Lenwood S. Heath. The pmu placement problem. *SIAM Journal on Discrete Mathematics*, 19:744–761, 1 2005.
- [9] Gerard Jennhwa Chang, Paul Dorbec, Mickael Montassier, and André Raspaud. Generalized power domination of graphs. *Discrete Applied Mathematics*, 160:1691–1698, 8 2012.
- [10] Changjie Cheng, Changhong Lu, and Yu Zhou. The k-power domination problem in weighted trees. *Theoretical Computer Science*, 809:231–238, 2 2020.
- [11] Nathaniel Dean, Alexandra Ilic, Ignacio Ramirez, Jian Shen, and Kevin Tian. On the power dominating sets of hypercubes. pages 488–491. IEEE, 8 2011.
- [12] Paul Dorbec, Antonio González, and Claire Pennarun. Power domination in maximal planar graphs. *Discrete Mathematics & Theoretical Computer Science*, 21:1–24, 6 2019.
- [13] Paul Dorbec, Michael A. Henning, Christian Löwenstein, Mickael Montassier, and André Raspaud. Generalized power domination in regular graphs. *SIAM Journal on Discrete Mathematics*, 27:1559–1574, 1 2013.
- [14] Paul Dorbec and Sandi Klavžar. Generalized power domination: Propagation radius and sierpiński graphs. *Acta Applicandae Mathematicae*, 134:75–86, 12 2014.
- [15] Paul Dorbec, Michel Mollard, Sandi Klavžar, and Simon Špacapan. Power domination in product graphs. *SIAM Journal on Discrete Mathematics*, 22:554–567, 1 2008.
- [16] Michael Dorfling and Michael A. Henning. A note on power domination in grid graphs. *Discrete Applied Mathematics*, 154:1023–1027, 4 2006.
- [17] Daniela Ferrero, Leslie Hogben, Franklin H. J. Kenter, and Michael Young. Note on power propagation time and lower bounds for the power domination number. *Journal of Combinatorial Optimization*, 34:736–741, 10 2017.
- [18] Jiong Guo, Rolf Niedermeier, and Daniel Raible. Improved algorithms and complexity results for power domination in graphs. *Algorithmica*, 52:177–202, 10 2008.

- [19] Teresa W. Haynes, Sandra M. Hedetniemi, Stephen T. Hedetniemi, and Michael A. Henning. Domination in graphs applied to electric power networks. *SIAM Journal on Discrete Mathematics*, 15:519–529, 1 2002.
- [20] Andreas M. Hinz, Seethu Varghese, and Ambat Vijayakumar. Power domination in knödel graphs and hanoi graphs. *Discussiones Mathematicae Graph Theory*, 38:63, 2018.
- [21] John J. Perry IV, Jason A. Perman, and Michael J. Zaworotko. Design and synthesis of metal–organic frameworks using metal–organic polyhedra as supermolecular building blocks. *Chemical Society Reviews*, 38:1400, 2009.
- [22] Liying Kang, Erfang Shan, and Min Zhao. Power domination in the generalized Petersen graphs. *Discussiones Mathematicae Graph Theory*, 40:695–712, 2020.
- [23] K.M. Koh and K.W. Soh. On the power domination number of the cartesian product of graphs. *AKCE International Journal of Graphs and Combinatorics*, 16:253–257, 12 2019.
- [24] Jyhmin Kuo and Wei-Lun Wu. Power domination in generalized undirected de Bruijn graphs and Kautz graphs. *Discrete Mathematics, Algorithms and Applications*, 07:1550003, 3 2015.
- [25] Meng-Yuan Li and Slavi C. Sevcik. Mellitate-based coordination polymers with a recurring motif: controlling dimensionality with secondary ligands. *CrystEngComm*, 15:5107, 2013.
- [26] Chung-Shou Liao and D. T. Lee. Power domination in circular-arc graphs. *Algorithmica*, 65:443–466, 2 2013.
- [27] Jia-Bao Liu, Muhammad Kashif Shafiq, Haidar Ali, Asim Naseem, Nayab Maryam, and Syed Sheraz Asghar. Topological indices of m th chain silicate graphs. *Mathematics*, 7:42, 1 2019.
- [28] Changhong Lu, Rui Mao, and Bing Wang. Power domination in regular claw-free graphs. *Discrete Applied Mathematics*, 284:401–415, 9 2020.
- [29] R. Sundara Rajan, J. Anitha, and Indra Rajasingh. 2-power domination in certain interconnection networks. *Procedia Computer Science*, 57:738–744, 2015.
- [30] Sudeep Stephen, Bharati Rajan, Joe Ryan, Cyriac Grigorious, and Albert William. Power domination in certain chemical structures. *Journal of Discrete Algorithms*, 33:10–18, 7 2015.

- [31] Chao Wang, Lei Chen, and Changhong Lu. k -power domination in block graphs. *Journal of Combinatorial Optimization*, 31:865–873, 2 2016.
- [32] S. Wilson. Power domination on permutation graphs. *Discrete Applied Mathematics*, 262:169–178, 6 2019.
- [33] Guangjun Xu and Liying Kang. On the power domination number of the generalized Petersen graphs. *Journal of Combinatorial Optimization*, 22:282–291, 8 2011.
- [34] Guangjun Xu, Liying Kang, Erfang Shan, and Min Zhao. Power domination in block graphs. *Theoretical Computer Science*, 359:299–305, 8 2006.
- [35] I Yuliana, Dafik, I H Agustin, and D A R Wardani. On the power domination number of corona product and join graphs. *Journal of Physics: Conference Series*, 1211:012020, 4 2019.
- [36] Min Zhao, Liying Kang, and Gerard J. Chang. Power domination in graphs. *Discrete Mathematics*, 306:1812–1816, 8 2006.
- [37] Min Zhao and Liying Kang. Power domination in planar graphs with small diameter. *Journal of Shanghai University (English Edition)*, 11:218–222, 6 2007.